

AD-A038 946

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER
ON AN EXTREMAL PROBLEM. (U)

F/6 12/1

JAN 77 P 6 NEVAI
MRC-TSR-1712

DAA629-75-C-0024
NL

UNCLASSIFIED

| OF |
AD
A038946



END

DATE
FILMED
5-77

AD A 038946

MRC Technical Summary Report #1712

ON AN EXTREMAL PROBLEM

Paul G. Nevai

12

Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53706

January 1977

(Received November 26, 1976)



Lee 1473

Approved for public release
Distribution unlimited

DDC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D. C.
20550

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

ON AN EXTREMAL PROBLEM

Paul G. Nevai

Technical Summary Report #1712
January 1977

ABSTRACT

Let $X = (x_1, x_2, \dots, x_N)$, $f: \mathbb{R} \rightarrow \mathbb{C}$ and let \mathbb{P}_n be the class of polynomials of degree at most n . The generalized Christoffel function Λ_n corresponding to the measure $d\alpha$ is defined by

$$\Lambda_n(X; f, N, d\alpha) = \min_{\substack{\pi \in \mathbb{P}_{n-1} \\ \pi(x_i) = f(x_i) \\ i=1, 2, \dots, N}} \int_{-\infty}^{\infty} |\pi(t)|^2 d\alpha(t).$$

It is shown that if α satisfies some rather weak conditions then

$\lim_{n \rightarrow \infty} n \Lambda_n(X; f, N, d\alpha)$ exists and the limit is also evaluated.

AMS (MOS) Subject Classification: 42A52

Key Words: Orthogonal polynomials, Christoffel functions

Work Unit Number 6 (Spline Functions and Approximation Theory)

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	Brit Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SE. 104
A	

ON AN EXTREMAL PROBLEM

Paul G. Nevai

The purpose of the present note is to investigate the asymptotic behavior of the functions $\Lambda_n : \mathbb{R}^N \rightarrow \mathbb{R}$ defined by

$$\Lambda_n(X; f, N, d\alpha) = \min_{\substack{\pi \in \mathbb{P}_{n-1} \\ \pi(x_i) = f(x_i) \\ i=1, 2, \dots, N}} \int_{-\infty}^{\infty} |\pi(t)|^2 d\alpha(t).$$

Here $X = (x_1, x_2, \dots, x_N)$, $f : \mathbb{R} \rightarrow \mathbb{C}$ is a fixed and almost everywhere finite function, \mathbb{P}_n is the set of all polynomials π of degree at most n and α is a weight function, that is α is nondecreasing on \mathbb{R} , it has infinitely many points of increase and every polynomial π belongs to $L_{d\alpha}^2$. Therefore Λ_n is defined and finite for almost every $X \in \mathbb{R}^N$.

Estimates for Λ_n lead to several results in probability theory, statistics and in the theory of orthogonal polynomials. (See [1], [2] and [4].) In fact, it is not hard to explicitly compute Λ_n ([3]) but the formula for Λ_n is so complicated that it cannot be used to estimate Λ_n when α is not nice. It will be shown that $\lim_{n \rightarrow \infty} n \Lambda_n(X; f, N, d\alpha)$ exists under rather weak assumptions on α and the corresponding limit will also be calculated.

Let $\{p_n(d\alpha)\}_{n=0}^{\infty}$ be the system of polynomials which is orthogonal with respect to $d\alpha$, that is $p_n(d\alpha, x) = \gamma_n x^n + \dots$ with $\gamma_n > 0$ and

$$\int_{-\infty}^{\infty} p_n(d\alpha, t) p_m(d\alpha, t) d\alpha(t) = \delta_{nm}.$$

Let M denote the class of those weights α for which

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} t p_n^2(d\alpha, t) d\alpha(t) = 0$$

and

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} t p_{n-1}(d\alpha, t) p_n(d\alpha, t) d\alpha(t) = \frac{1}{2}.$$

Let us remark that M contains many weights. If, for instance, $\text{supp}(d\alpha) = [-1, 1]$ and $\log \alpha'(\cos \theta) \in L^1$ then $\alpha \in M$ ([5]).

Furthermore, if α is absolutely continuous, $\text{supp}(d\alpha) = [-1, 1]$ and $\alpha'(t) = \varphi(t) \exp\{-(1-t^2)^{-\frac{1}{2}}\}$ where $0 < c \leq \varphi(t)$ for $-1 \leq t \leq 1$ and φ is Riemann integrable then also $\alpha \in M$ ([4]). Another example is the Pollaczek weight ([5]). In the following Δ denotes an interval.

It is known that if $\alpha \in M$ then $\Delta \subset \text{supp}(d\alpha)$ iff $\Delta \subset [-1, 1]$ ([4]).

THEOREM. Let $\alpha \in M$, $\Delta \subset \text{supp}(d\alpha)$, $1/\alpha' \in L^1(\Delta)$. Let exist a sequence $\{\varepsilon_k\}_{k=1}^{\infty}$ with $\varepsilon_k \geq 0$, $\varepsilon_k \searrow 0$ such that for every fixed k the function Ψ_k , which is defined by $\Psi_k(t) = [(1 - \varepsilon_k)^2 - t^2]^{-\frac{1}{2}} \log \alpha'(t)$, belongs to $L^1(-1 + \varepsilon_k, 1 - \varepsilon_k)$. Then for almost every $X \in \Delta^N$

$$\lim_{n \rightarrow \infty} n \Lambda_n(X; f, N, d\alpha) = \pi \sum_{i=1}^N |f(x_i)|^2 \alpha'(x_i) \sqrt{1 - x_i^2}.$$

Proof. For $N = 1$ the Theorem has been proved in [4]. Now let $N > 1$.

If $\pi \in \mathbb{P}_{n-1}$ then π can be expressed as

$$\pi(x) = \int_{-\infty}^{\infty} \pi(t) K_n(d\alpha, x, t) d\alpha(t)$$

where

$$K_n(d\alpha, x, t) = \sum_{k=0}^{n-1} p_k(d\alpha, x) p_k(d\alpha, t).$$

If X is given and $\prod_{i < j} (x_i - x_j) \neq 0$ then we can write

$$\begin{aligned} \sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) &= \\ &= \int_{-\infty}^{\infty} \pi(t) \left[\sum_{i=1}^N \text{sign } \pi(x_i)^2 \pi(x_i) \Lambda_n(x_i, 1, 1, d\alpha) K_n(d\alpha, x_i, t) \right] d\alpha(t). \end{aligned}$$

Using Cauchy's inequality, orthogonality relations and the well known fact that

$$(1) \quad K_n(d\alpha, x, x) = \Lambda_n(x; 1, 1, d\alpha)$$

we obtain

$$\begin{aligned} \left[\sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) \right]^2 &\leq \\ &\leq \int_{-\infty}^{\infty} |\pi(t)|^2 d\alpha(t) \cdot \left[\sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) + \right. \\ &\quad \left. + 2 \sum_{j < i} |\pi(x_i)| |\pi(x_j)| \Lambda_n(x_i; 1, 1, d\alpha) \Lambda_n(x_j; 1, 1, d\alpha) |K_n(d\alpha, x_i, x_j)| \right]. \end{aligned}$$

If $\alpha \in M$ then clearly $\text{supp}(d\alpha)$ is compact. Therefore by the Christoffel-Darboux formula there exists a number C depending on X and $\text{supp}(d\alpha)$ such that

$$|K_n(d\alpha, x_i, x_j)| \leq C[|p_n(d\alpha, x_i)p_{n-1}(d\alpha, x_j)| + |p_{n-1}(d\alpha, x_i)p_n(d\alpha, x_j)|].$$

Hence

$$\begin{aligned} & \left[\sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) \right]^2 \leq \\ & \leq \int_{-\infty}^{\infty} |\pi(t)|^2 d\alpha(t) \left[\sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) + \right. \\ & \quad + 4C \sum_{i=1}^N |\pi(x_i)| |p_n(d\alpha, x_i)| \Lambda_n(x_i; 1, 1, d\alpha) \cdot \\ & \quad \cdot \sum_{i=1}^N |\pi(x_i)| |p_{n-1}(d\alpha, x_i)| \Lambda_n(x_i; 1, 1, d\alpha) \left. \right]. \end{aligned}$$

Formula (1) implies that $p_{n-1}^2(d\alpha, x_i) \Lambda_n(x_i; 1, 1, d\alpha) \leq 1$. Consequently by Cauchy's inequality

$$\begin{aligned} & \left[\sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) \right]^2 \leq \\ & \leq \int_{-\infty}^{\infty} |\pi(t)|^2 d\alpha(t) \sum_{i=1}^N \Lambda_n(x_i; |\pi(x_i)|, 1, d\alpha) \cdot \\ & \quad \cdot \{1 + 4C\sqrt{N} \left[\sum_{i=1}^N \Lambda_n(x_i; |p_n(d\alpha, x_i)|, 1, d\alpha) \right]^{\frac{1}{2}}\}. \end{aligned}$$

This inequality holds for each $X \in \mathbb{R}^N$ such that $\prod_{i < j} (x_i - x_j) \neq 0$.

It has been proved in [4] that if $\alpha \in M$ then for every $x \in \text{supp}(d\alpha)$

$$(2) \quad \lim_{n \rightarrow \infty} \Lambda_n(x; |p_n(d\alpha, x)|, 1, d\alpha) = 0.$$

Consequently if $\alpha \in M$ then for almost every $X \in [\text{supp}(d\alpha)]^N$

$$(3) \quad \sum_{i=1}^N \Lambda_n(x_i; |f(x_i)|, 1, d\alpha) \leq [1 + \sigma(1)] \Lambda_n(X, f, N, d\alpha)$$

where $\lim_{n \rightarrow \infty} \sigma(1) = 0$ for almost every $X \in [\text{supp}(d\alpha)]^N$. Our next

aim is to establish the converse inequality. For a given X let π^X be defined by

$$\pi^X(t) = \sum_{i=1}^N f(x_i) \Lambda_{n-N+1}(x_i; 1, 1, d\alpha) K_{n-N+1}(d\alpha, t, x_i) \ell_i(t)$$

where $\{\ell_i\}_{i=1}^N$ are the fundamental polynomials of Lagrange interpolation corresponding to $\{x_i\}_{i=1}^N$. Clearly $\pi^X \in \mathbb{P}_{n-1}$ for almost every $X \in \mathbb{R}^N$.

Furthermore, if $\pi^X \in \mathbb{P}_{n-1}$ then $\pi^X(x_i) = f(x_i)$ for $i = 1, 2, \dots, N$.

Let us compute the $L_{d\alpha}^2$ norm of π^X . Introducing the notation

$$G_n(d\alpha, g, x) = \Lambda_n(x; 1, 1, d\alpha) \int_{-\infty}^{\infty} g(t) K_n^2(d\alpha, x, t) d\alpha(t)$$

we have

$$(4) \quad \int_{-\infty}^{\infty} |\pi^X(t)|^2 d\alpha(t) = \sum_{i=1}^N |f(x_i)|^2 \Lambda_{n-N+1}(x_i; 1, 1, d\alpha) G_{n-N+1}(d\alpha, \ell_i^2, x_i) \\ + 2 \sum_{i < j} \text{Re}[f(x_i) \overline{f(x_j)}] \Lambda_{n-N+1}(x_i; 1, 1, d\alpha) \Lambda_{n-N+1}(x_j; 1, 1, d\alpha) \cdot \\ \cdot \int_{-\infty}^{\infty} \ell_i(t) \ell_j(t) K_{n-N+1}(d\alpha, t, x_i) K_{n-N+1}(d\alpha, t, x_j) d\alpha(t).$$

If $\alpha \in M$, g is continuous on $\text{supp}(d\alpha)$ and $x \in \text{supp}(d\alpha)$ then

$$\lim_{n \rightarrow \infty} G_n(d\alpha, g, x) = g(x)$$

(see [4]), in particular, if X is such that $\prod_{i < j} (x_i - x_j) \neq 0$ then

$$\lim_{n \rightarrow \infty} G_n(d\alpha, \ell_i^2, x_i) = \ell_i^2(x_i) = 1.$$

Let

$$I_{ij} = \int_{-\infty}^{\infty} \ell_i(t) \ell_j(t) K_{n-N+1}(d\alpha, t, x_i) K_{n-N+1}(d\alpha, t, x_j) d\alpha(t).$$

If $\prod_{k < \ell} (x_k - x_\ell) \neq 0$ then $\ell_i \ell_j \in \mathbb{P}_{2N-2}$. By a direct calculation we obtain that

$$\begin{aligned} I_{ij} &= \int_{-\infty}^{\infty} \ell_i(t) \ell_j(t) \left[K_{n-3N+3}(d\alpha, x_i, t) + \sum_{k=n-3N+3}^{n-N} p_k(d\alpha, x_i) p_k(d\alpha, t) \right] \cdot \\ &\quad \cdot K_{n-N+1}(d\alpha, x_j, t) d\alpha(t) = \ell_i(x_j) \ell_j(x_j) K_{n-3N+3}(d\alpha, x_i, x_j) + \\ &\quad + \sum_{k=n-3N+3}^{n-N} p_k(d\alpha, x_i) \sum_{\ell=k-2N+2}^{n-N} p_\ell(d\alpha, x_j) \cdot \\ &\quad \cdot \int_{-\infty}^{\infty} \ell_i(t) \ell_j(t) p_k(d\alpha, t) p_\ell(d\alpha, t) d\alpha(t). \end{aligned}$$

Since $\ell_i(x_j) = 0$ we get

$$(5) \quad |I_{ij}| \leq C_1 \sum_{k=n-5N}^{n-N} |p_k(d\alpha, x_i)| \sum_{\ell=n-5N}^{n-N} |p_\ell(d\alpha, x_j)|$$

where C_1 depends on X and $\text{supp}(d\alpha)$. Using the recurrence formula

which the orthogonal polynomials satisfy it can easily be seen that

for $\alpha \in M$ (5) implies

$$|I_{ij}| \leq C_2 \sum_{k=n-N-1}^{n-N} |p_k(d\alpha, x_i)| \sum_{\ell=n-N-1}^{n-N} |p_\ell(d\alpha, x_j)|$$

where C_2 is independent of n . Applying now (1) and (2) the second sum on the right hand side of (4) can easily be estimated. We get

$$\int_{-\infty}^{\infty} |\pi^X(t)|^2 d\alpha(t) = [1 + \sigma(1)] \sum_{i=1}^N |f(x_i)|^2 \Lambda_{n-N+1}(x_i; 1, 1, d\alpha)$$

for almost every $X \in [\text{supp}(d\alpha)]^N$. It has been proved in [4] that if $\alpha \in M$ and $x \in \text{supp}(d\alpha)$ then

$$\lim_{n \rightarrow \infty} \Lambda_{n-N+1}(x; 1, 1, d\alpha) \Lambda_n(x; 1, 1, d\alpha)^{-1} = 1$$

for every fixed N . Thus

$$(6) \quad \Lambda_n(X; f, N, d\alpha) \leq [1 + \sigma(1)] \sum_{i=1}^N \Lambda_n(x_i; |f(x_i)|, 1, d\alpha)$$

for almost every $X \in [\text{supp}(d\alpha)]^N$. The Theorem follows now from estimates (3) and (6).

REFERENCES

- [1] Freud, G., Orthogonal Polynomials, Pergamon Press, New York, 1971.
- [2] Grenander, U. and Szegő, G., Toeplitz Forms and Their Applications, University of California Press, Berkeley, 1958.
- [3] Grenander, U. and Rosenblatt, M., An extension of a theorem of G. Szegő and its application to the study of stochastic processes, Trans. Amer. Math. Soc. 76 (1954), 112-126.
- [4] Nevai, P., Orthogonal Polynomials (to appear).
- [5] Szegő, G., Orthogonal Polynomials, Amer. Math. Soc., Providence, Rh. I., 1967.

Department of Mathematics and
Mathematics Research Center
University of Wisconsin
Madison, Wisconsin 53706

and

Department of Mathematics
The Ohio State University
Columbus, Ohio 43210

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1712	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9) Technical
4. TITLE (and Subtitle) (6) ON AN EXTREMAL PROBLEM.		5. TYPE OF REPORT & PERIOD COVERED Summary Report, - no specific reporting period
7. AUTHOR(s) (10) Paul G. Nevai		6. PERFORMING ORG. REPORT NUMBER (15)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ✓ NSF-MCS-75-06687 #3
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 6 (Spline Functions and Approximation Theory)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 11p.		12. REPORT DATE (11) January 1977
		13. NUMBER OF PAGES 8
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) (14) MRC-TSR-1712 Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Orthogonal polynomials Christoffel functions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Asymptotic estimate is given for a generalization of Christoffel functions.		